


AD-A185 611 PEAKEDNESS OF WEIGHTED AVERAGES OF JOINTLY DISTRIBUTED 1/1
RANDOM VARIABLES. (U) FLORIDA STATE UNIV TALLAHASSEE
DEPT OF STATISTICS W CHAN ET AL. AUG 87
UNCLASSIFIED FSU-STATISTICS-TR-M712R AFOSR-TR-87-1574 F/G 12/3 NL





AD-A185 611

DTIC FILE COPY
TATION PAGE

RESTRICTIVE MARKINGS

2

1a. REPORT SECURITY

UNCLASSIFIED

2a. SECURITY CLASSIFICATION AUTHORITY

NA

2b. DECLASSIFICATION/DOWNGRADING SCHEDULE

NA

3. DISTRIBUTION/AVAILABILITY OF REPORT

Approved for Public Release; Distribution Unlimited.

4. PERFORMING ORGANIZATION REPORT NUMBER(S)

FSU TECHNICAL REPORT NO M712R

5. MONITORING ORGANIZATION REPORT NUMBER(S)

AFOSR-TR-87-1574

6a. NAME OF PERFORMING ORGANIZATION
FLORIDA STATE UNIVERSITY6b. OFFICE SYMBOL
(If applicable)7a. NAME OF MONITORING ORGANIZATION
AFOSR/NM

6c. ADDRESS (City, State and ZIP Code)

Department of Statistics
Tallahassee, FL 32306-3033

7b. ADDRESS (City, State and ZIP Code)

Bldg. 410
Bolling AFB, DC 20332-64488a. NAME OF FUNDING/SPONSORING
ORGANIZATION
AFOSR8b. OFFICE SYMBOL
(If applicable)
NM

9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER

F49620-82-K-0007

8c. ADDRESS (City, State and ZIP Code)

Bldg. 410
Bolling AFB, DC

10. SOURCE OF FUNDING NOS.

PROGRAM
ELEMENT NO.
6.1102FPROJECT
NO.
2304TASK
NO.
A-5WORK UNIT
NO.

11. TITLE (Include Security Classification)

PEAKEDNESS OF WEIGHTED AVERAGES OR JOINTLY DISTRIBUTED
RANDOM VARIABLES

12. PERSONAL AUTHOR(S)

Wai Chan, Dong Ho Park, and Frank Proschan

13a. TYPE OF REPORT

Journal
Technical Report

13b. TIME COVERED

FROM TO

14. DATE OF REPORT (Yr., Mo., Day)

August 1987

15. PAGE COUNT

1-6

16. SUPPLEMENTARY NOTATION

17. COSATI CODES

FIELD	GROUP	SUB. GR.

18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)

Key Words - Peakedness, convex combination, majorization,
Schur-concave density, Cauchy distribution.

19. ABSTRACT (Continue on reverse if necessary and identify by block number)

Sub 2
This note extends the Proschan (1965) result on peakedness comparison for convex combinations of i.i.d. random variables from a PF_2 density. Now the underlying random variables are jointly distributed from a Schur-concave density. The result permits a more refined description of convergence in the Law of Large Numbers.

Cauchy distribution
convergence

DTIC

ELECTE

OCT 29 1987

20. DISTRIBUTION/AVAILABILITY OF ABSTRACT

UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT. ☐ DTIC USERS ☐

21. ABSTRACT SECURITY CLASSIFICATION

UNCLASSIFIED

22a. NAME OF RESPONSIBLE INDIVIDUAL

Frank Proschan

May Woodruff

22b. TELEPHONE NUMBER
(Include Area Code)

(904) 644-3218

22c. OFFICE SYMBOL

AFOSR/NM

REVISION

AFOSR-TR- 87-1574

**PEAKEDNESS OF WEIGHTED AVERAGES OF
JOINTLY DISTRIBUTED RANDOM VARIABLES**

by

Wai Chan¹, Dong Ho Park², and Frank Proschan³

**FSU Technical Report No. M 712R
AFOSR Technical Report No. 85-184R**

August, 1987

**¹Ohio State University
Department of Statistics
Columbus, Ohio 43210**

and

**²University of Nebraska
Department of Statistics
Lincoln, Nebraska 68588**

and

**³Florida State University
Department of Statistics
Tallahassee, Florida 32306**

Research sponsored by the Air Force Office of Scientific Research under Contract Number AFOSR 82-K-0007. The U.S. Government is authorized to reproduce and distribute reprints for Governmental purposes notwithstanding any copyright notation thereon.

Key Words: Peakedness, convex combination, majorization, Schur-concave density, Cauchy distribution.

AMS (1981) subject classification No. 60E15.

**PEAKEDNESS OF WEIGHTED AVERAGES OF
JOINTLY DISTRIBUTED RANDOM VARIABLES**

by

Wai Chan, Dong Ho Park, and Frank Proschan

ABSTRACT

This note extends the Proschan (1965) result on peakedness comparison for convex combinations of i.i.d. random variables from a PF_2 density. Now the underlying random variables are jointly distributed from a Schur-concave density. The result permits a more refined description of convergence in the Law of Large Numbers.



Accession For	
NTIS CRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution	
Availability	
Dist	Availability for Special
A-1	

1. Introduction

Proschan (1965) shows that:

1.1 Theorem. Let f be PF_2 , $f(t) = f(-t)$ for all t , X_1, \dots, X_n independently distributed with density f , $\underline{p} \stackrel{m}{\geq} \underline{p}'$, $\underline{p}, \underline{p}'$ not identical, $\sum_1^n p_i = 1 = \sum_1^n p'_i$. Then $\sum_1^n p_i X_i$ is strictly more peaked than $\sum_1^n p'_i X_i$.

(Definitions of majorization ($\underline{p} \stackrel{m}{\geq} \underline{p}'$), PF_2 density, and peakedness are presented in Section 2.)

The Law of Large Numbers asserts that the average of a random sample converges to the population mean under certain conditions. Roughly speaking, Theorem 1.1 states that a weighted average of i.i.d. random variables converges more rapidly in the case in which weights are close together as compared with the case in which the weights are diverse.

In the present note, we extend the basic univariate result to the multivariate situation in which the underlying random variables have a joint Schur-concave density. Theorem 2.4 presents the precise statement of the multivariate extension.

2. Peakedness comparisons

The theory of majorization is exploited in this section to obtain more general versions of the result of Proschan (1965). We begin with some definitions. The definition of peakedness was given by Birnbaum (1948).

Definition 2.1. Let X and Y be real valued random variables and a and b real constants. We say that X is more peaked about a than Y about b if

$$P(|X-a| \geq t) \leq P(|Y-b| \geq t)$$

for all $t \geq 0$. In the case $a = 0 = b$, we simply say that X is more peaked than Y .

Next we define the ordering of majorization among vectors.

Definition 2.2. Let $a_1 \geq \dots \geq a_n$ and $b_1 \geq \dots \geq b_n$ be decreasing rearrangements of the components of the vectors a and b . We say that a majorizes b (written $a \overset{m}{\geq} b$) if

$$\sum_{i=1}^n a_i = \sum_{i=1}^n b_i$$

and

$$\sum_{i=1}^k a_i \geq \sum_{i=1}^k b_i \quad \text{for } k = 1, \dots, n-1.$$

Definition 2.3. A real valued function f defined on R^n is said to be a Schur-concave function if $f(a) \leq f(b)$ whenever $a \geq b$.

A nonnegative function f on $(-\infty, \infty)$ is called a Pólya frequency function of order 2 (PF_2) if $\log f$ is concave. If f is a PF_2 function then $\phi(x) = \prod f(x_i)$ is Schur-concave. Thus the random vector $x = (X_1, \dots, X_n)$ has a Schur-concave density under the conditions of Theorem 1.1. A function f defined on R^n is said to be sign-invariant if $f(x_1, \dots, x_n) = f(|x_1|, \dots, |x_n|)$. In the following theorem, we give a peakedness comparison for random variables with a sign-invariant Schur-concave density.

Theorem 2.4. Suppose the random vector $X = (X_1, \dots, X_n)$ has a sign invariant Schur-concave density. Then for all $t \geq 0$,

$$\psi(a_1, \dots, a_n) = P(\sum a_i X_i \leq t)$$


is a Schur-concave function of $\underline{a} = (a_1, \dots, a_n)$, $a_i \geq 0$ for all i . Equivalently, $\sum b_i X_i$ is more peaked than $\sum a_i X_i$ whenever $\underline{a} \overset{m}{\geq} \underline{b}$.

Proof.

Without loss of generality, we may assume that $\sum a_i = 1$. We first consider the case $n = 2$.

Let $\underline{a} = (a_1, a_2)$, $\underline{b} = (b_1, b_2)$, $\underline{a} \stackrel{m}{\geq} \underline{b}$. Since X_1, X_2 are exchangeable, we may further assume that

$a_1 > b_1 \geq 1/2 \geq b_2 > a_2$. To show that $P(a_1 X_1 + a_2 X_2 \leq t) \leq P(b_1 X_1 + b_2 X_2 \leq t)$ for $t \geq 0$, consider the following diagram:



Since $a_1 > b_1 \geq 1/2$, both lines intersect the x_1 -axis in the interval $[t, 2t]$ and they intersect the 45 degree line at the point (t, t) ($a_1 + a_2 = b_1 + b_2 = 1$). We must show that $P(E) \leq P(F)$. Now reflect E across the 45 degree line to form the wedge E' . Then $P(E) = P(E')$ because the joint density f is invariant under permutation. For $k \geq 0$, the line $x_1 - x_2 = k$ intersects E' at the line segment joining $(t+b_1k, t-b_2k)$ and $(t+a_1k, t-a_2k)$, and it intersects F at the line segment joining $(t+a_2k, t-a_1k)$ and $(t+b_2k, t-b_1k)$. Note that both segments are of equal length. But f sign-invariant and Schur-concave implies that

$$f(t+b_1k, t-b_2k) = f(t+b_1k, b_2k-t)$$

$$\leq f(t+b_2k, b_1k-t)$$

$$= f(t+b_2k, t-b_1k).$$

This last fact then clearly implies that $P(E') \leq P(F)$ by conditioning on $X_1 - X_2$.

The result for $n \geq 3$ now follows since

$$P(\sum a_i X_i \leq t)$$

$$= E [P(a_1 X_1 + a_2 X_2 \leq t - \sum_{i=3}^n a_i X_i \mid X_3, \dots, X_n)]$$

and the conditional density $f(x_1, x_2 \mid x_3, \dots, x_n)$ is also Schur-concave and sign invariant.

For an example of a Schur-concave density function that is also sign-invariant, consider the multivariate Cauchy density:

$$f(x_1, \dots, x_n) = \pi^{-(n+1)/2} \Gamma((n+1)/2) (1 + \sum_{i=1}^n x_i^2)^{-(n+1)/2}$$

The following result is an immediate consequence of Theorem 2.4.

Corollary 2.5. Let X_1, \dots, X_n be random variables with joint Schur-concave sign-invariant density f . Then $\frac{1}{k} \sum_{i=1}^k X_i$ is increasing in peakedness as k increases from 1 to n .

Proof.

Let $\underline{a}_1 = (1, 0, \dots, 0)$, $\underline{a}_2 = (\frac{1}{2}, \frac{1}{2}, 0, \dots, 0)$, ..., and $\underline{a}_n = (\frac{1}{n}, \dots, \frac{1}{n})$, where each vector contains n components. Then $\underline{a}_1 \succeq^m \underline{a}_2 \succeq^m \dots \succeq^m \underline{a}_n$. The result follows from Theorem 2.4.

Suppose $\underline{X} = (X_1, \dots, X_n)$ and $\underline{Y} = (Y_1, \dots, Y_n)$ are independently distributed with respective densities f and g where both f and g are Schur-concave and sign invariant. Then Theorem 2.4 implies that $\sum b_i (X_i + Y_i)$ is more peaked than $\sum a_i (X_i + Y_i)$ whenever $\underline{a} \succeq^m \underline{b}$. This is true because the convolution of Schur-concave functions is Schur-concave. However, if Y_1, \dots, Y_n are i.i.d. Cauchy, then the joint density g given by

$$(2.1) \quad g(y_1, \dots, y_n) = \left(\frac{a}{\pi}\right)^n \prod_{i=1}^n (1 + a^2 y_i^2)^{-1}, \quad a > 0,$$

is not Schur-concave. In Theorem 2.6 below, we show that $\sum b_i (X_i + Y_i)$ is more peaked than $\sum a_i (X_i + Y_i)$ whenever $\underline{a} \succeq^m \underline{b}$. This result identifies a different class of densities for which the conclusion of Theorem 2.4 holds.

Theorem 2.6. Suppose that the random vector $\underline{X} = (X_1, \dots, X_n)$ has a sign-invariant Schur-concave density f . Let $\underline{Y} = Y_1, \dots, Y_n$ be i.i.d. Cauchy with joint density g as given in (2.1). Let \underline{X} and \underline{Y} be independent, and $\underline{a} \succeq \underline{b}$ where $a_i \geq 0, b_i \geq 0$ for all i and $1 = \sum_1^n a_i = \sum_1^n b_i$. Then $\sum_1^n b_i (X_i + Y_i)$ is more peaked than $\sum_1^n a_i (X_i + Y_i)$.

Proof.

Since f is sign invariant both $\sum_1^n a_i X_i$ and $\sum_1^n b_i X_i$ are symmetric random variables. We use the

fact that $\sum_1^n a_i Y_i, \sum_1^n b_i Y_i$ have the same distribution as does Y_1 . The result now follows

Theorem 2.4 and the Lemma of Birnbaum (1948) by noting that Y_1 has a symmetric and unimodal density.

References

- [1] Birnbaum, Z.W. (1948). On random variables with comparable peakedness.
Ann. Math. Statist. 19 76-81.
- [2] Proschan, F. (1965). Peakedness of distributions of convex combinations.
Ann. Math. Statist. 36 1703-1706.

END

12-87

DTIC